



Lecture (05) Electric Potential


By:

Dr. Ahmed ElShafee

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II


Electrostatic Potential Energy and Potential Difference

- The Coulomb (Electrostatic) force is a conservative force (A *conservative force* is a *force* with the property that the work done in moving a particle between two points is independent of the taken path.)
- A potential energy function can be defined for any conservative force, including Coulomb force



introduction

- We are used to voltage in our lives—a 12-volt car battery, 110 V or 220 V at home, 1.5 volt flashlight batteries, and so on.
- Voltage is the same as electric potential difference between two points.
- Electric potential is defined as the potential energy per unit charge.




Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

- As in mechanics, work is

$$w = f \times d$$

- Work done on the positive charge by moving it from A to B

$$w = E \times q \times d$$



- The work done by a conservative force equals the **negative** of the **change in potential energy**, ΔPE

$$\Delta PE(\Delta U) = -W = -fd = -qEd$$

- This equation
 - is valid only for the case of a **uniform** electric field
 - allows to introduce the concept of **electric potential**

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

- Electric potential difference is the work done to move a charge from a point A to a point B divided by the magnitude of the charge.

- Thus the SI units of electric potential

$$1v = 1 \frac{J}{C}$$

- In other words, 1 J of work is required to move a 1 C of charge between two points that are at potential difference of 1 V

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

- Electric potential
- The **potential difference** between points A and B, $V_B - V_A$, is defined as the change in potential energy (final minus initial value) of a charge, q, moved from A to B, divided by the charge
- Type equation here.


$$\Delta V = V_B - V_A = \frac{\Delta PE(\Delta U)}{q}$$

$$= \frac{U_b - U_a}{q} = -Ed$$

- Electric potential is a **scalar quantity**
- Electric potential difference is a measure of electric energy per unit charge

- Potential is often referred to as “voltage”

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II



- Units of electric field (N/C) can be expressed in terms of the units of potential (as volts per meter)


$$\Delta V = V_B - V_A = \frac{\Delta PE(\Delta U)}{q} = -Ed = -\frac{F}{q} d$$

$$\frac{\Delta V}{d} = -\frac{F}{q} \frac{1}{d} = -\frac{1}{d} \frac{J}{C} = -\frac{1}{m} \frac{J}{C}$$

- Because the positive tends to move in the direction of the electric field, work must be done on the charge to move it in the direction, opposite the field

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

- Thus,
 - A positive charge gains electric potential energy when it is moved in a direction opposite the electric field
 - A negative charge loses electrical potential energy when it moves in the direction opposite the electric field



Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

MCQ

- A proton and an electron are in a constant electric field created by oppositely charged plates. You release the proton from the positive side and the electron from the negative side. Which has the larger acceleration?


1. proton
2. electron
3. both feel the same acceleration
4. neither – there is no acceleration
5. they feel the same magnitude acceleration but opposite direction

Since $F = ma$ and the electron is much less massive than the proton, the electron experiences the larger acceleration.

MCQ

- A proton and an electron are in a constant electric field created by oppositely charged plates. You release the proton from the positive side and the electron from the negative side. Which has the larger acceleration?

1. proton
2. electron
3. both feel the same acceleration
4. neither – there is no acceleration
5. they feel the same magnitude acceleration but opposite direction




Analogy between gravitational and electrical potential energy

$$W = f d = E Q d$$


$$\Delta V = -E d$$

$$W = -Q \Delta v$$



Example 01


- Suppose a negative charge, such as an electron, is placed near the negative plate at point b, as shown here. If the electron is free to move, will its electric potential energy increase or decrease? How will the electric potential change?



13

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

- (a) Two rocks are at the same height.
- The larger rock has more potential energy.
- (b) Two charges have the same electric potential.
- The $2Q$ charge has more potential energy.



14

- The electron will move towards the positive plate if released, thereby increasing its kinetic energy.
- Its potential energy must therefore decrease. However, it is moving to a region of higher potential V ; the potential is determined only by the existing charge distribution and not by the point charge.

- Electrical sources such as batteries and generators supply a constant potential difference. Here are some typical potential differences, both natural and manufactured:

TABLE 23-1 Some Typical Potential Differences (Volts)

Source	Voltage (approx.)
Thundercloud to ground	10^8 V
High-voltage power line	10^5 – 10^6 V
Power supply for TV tube	10^4 V
Automobile ignition	10^4 V
Household outlet	10^2 V
Automobile battery	12 V
Flashlight battery	1.5 V
Resting potential across nerve membrane	10^{-1} V
Potential changes on skin (EKG and EEG)	10^{-4} V

15


Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

16

Dr. Ahmed ElShafee, ACU

Example 02

- Suppose an electron in a cathode ray tube is accelerated from rest through a potential difference $V_b - V_a = V_{ba} = +5000 \text{ V}$
- (a) What is the change in electric potential energy of the electron?
- (b) What is the speed of the electron ($m = 9.1 \times 10^{-31} \text{ kg}$) as a result of this acceleration?



Dr. Ahmed ElShafee, ACU : Spring 2015, Ph


14

MCQ

- Which requires the **most work**, to move a positive charge from **P** to points **1, 2, 3** or **4**? All points are the same distance from **P**.


- $P \rightarrow 1$
- $P \rightarrow 2$
- $P \rightarrow 3$
- $P \rightarrow 4$

- 5) all require the same amount of work



Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

14



- $\Delta U = F d = E q d$
 - But $\Delta V = E d$
 - $\therefore \Delta U = \Delta V \times q = 5000 \times -1.6 \times 10^{-19} = -8.0 \times 10^{-16} \text{ J}$

- The potential energy lost by the electron becomes kinetic energy K . From conservation of energy $\Delta K + \Delta U = 0$, so

$$\Delta K = -\Delta U$$

$$\frac{1}{2} m v^2 - 0 = -q(v_b - v_a) = -q v_{ba}$$

$$v = \sqrt{\frac{-2q v_{ba}}{m}} = \sqrt{\frac{-2 \times -1.6 \times 10^{-19} \times 5000}{9.1 \times 10^{-31}}} = 4.2 \times 10^7 \text{ m/s}$$

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II


14

MCQ

- Which requires the **most work**, to move a positive charge from **P** to points **1, 2, 3** or **4**? All points are the same distance from **P**.

- $P \rightarrow 1$
- $P \rightarrow 2$
- $P \rightarrow 3$
- $P \rightarrow 4$

- 5) all require the same amount of work




For **path #1**, you have to push the positive charge **against** the E field, which is **hard to do**. By contrast, path #4 is the **easiest**, since the field does all the work.

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

14

Relation between Electric Potential and Electric Field

- To find V_{ba} in a nonuniform electric field E , we integrate $E \cdot dl$ from point a to point b.
- The general relationship between a conservative force and potential energy
- $U_b - U_a = - \int_a^b f dl$
- Substituting the potential difference and the electric field:
- $v_{ba} = v_b - v_a = - \int_a^b E dl$



Y1

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II


- $\Delta PE (\Delta U) = -w = -f d$
- but $f = k \frac{q_1 \times q_2}{d^2}$
- $\Delta PE (\Delta U) = -k \frac{q_1 \times q_2}{d} = \frac{-q_1 \times q_2}{4 \pi \epsilon_0 d}$

Y1

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

- A simple special case is a uniform field. In Fig., for example, a path parallel to the electric field lines from point a at the positive plate to point b at the negative plate gives (since E and dl are in the same direction at each point),

$$v_{ba} = -E \int_a^b dl = -E d$$




Y1

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

Example 03

- Two parallel plates are charged to produce a potential difference of 50 V. If the separation between the plates is 0.050 m, calculate the magnitude of the electric field in the space between the plates.




Y1

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

Example 04


- $V = E d$
- $E = V/d = 50/0.05 = 1000 \text{ V/m}$.



٧٥

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

- Determine the potential at a distance r from the center of a uniformly charged conducting sphere of radius r_0 for
 - (a) $r > r_0$,
 - (b) $r = r_0$,
 - (c) $r < r_0$.
- The total charge on the sphere is Q .




٧٦

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

a)

- for $r > r_0$
- $E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$
- $v_b - v_a = -\int_{r_a}^{r_b} E dr$
- $= \frac{-Q}{4\pi\epsilon_0} \int_{r_a}^{r_b} \frac{dr}{r^2} = \frac{-Q}{4\pi\epsilon_0} \left(\frac{1}{r_b} - \frac{1}{r_a} \right)$
- set $r_a = r$ and $r_b = \infty$
- $v_{ba} = \frac{Q}{4\pi\epsilon_0 r}$
- $v = \frac{Q}{4\pi\epsilon_0 r}$



٧٧

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

b) $r = r_0 \rightarrow$ conductor has same potential over its surface

$$v = \frac{Q}{4\pi\epsilon_0 r_0}$$

c) $r < r_0 \rightarrow$ conductor has same potential

$$v = \frac{Q}{4\pi\epsilon_0 r_0}$$


٧٨

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

- The previous example gives the electric potential as a function of distance from the surface of a charged conducting sphere, which is plotted here, and compared with the electric field:

$$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$$

$$V = \frac{Q}{4\pi\epsilon_0 r}$$





Y4

Dr. Ahmed ElShafee, ACU : Spring 201



Y4

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II



$$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$$

$$V = \frac{Q}{4\pi\epsilon_0 r_0}$$



Y4

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

Y4

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II


Electric Potential Due to Point Charges

- To find the electric potential due to a point charge, we integrate the field along a field line

$$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$$

$$V_b - V_a = - \int_{r_a}^{r_b} E dl$$

$$= \frac{-Q}{4\pi\epsilon_0} \int_{r_a}^{r_b} \frac{dr}{r^2} = \frac{-Q}{4\pi\epsilon_0} \left(\frac{1}{r_b} - \frac{1}{r_a} \right)$$



Electrostatic Potential Energy; between two charges



- The potential energy of a charge in an electric potential is

$$U = qV.$$

- To find the electric potential energy of two charges, imagine bringing each in from infinitely far away.
- The first one takes no work, as there is no field.
- To bring in the second one, we must do work due to the field of the first one; this means the potential energy of the pair is:

$$U = q_2V = k \frac{q_1 q_2}{r_{12}}$$

٧٤



Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

٧٧

Example 05

- Example: Work required to bring two positive charges close together.
- What minimum work must be done by an external force to bring a proton $q = 1.60 \times 10^{-19}$ C from a great distance away (take $r = \infty$) to a point 1.60×10^{-15} m from another proton?

$q = 1.60 \times 10^{-19}$

1.60×10^{-15} m

$F \infty$
 $q = 1.60 \times 10^{-19}$

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

٧٥

1.60×10^{-15} m

$q = 1.60 \times 10^{-19}$

$F \infty$
 $q = 1.60 \times 10^{-19}$

- The work is equal to the change in potential energy
- $\Delta W = -q \Delta v = -q(v_b - v_a) = -q \left(\frac{kq}{r_b} - \frac{kq}{r_a} \right)$
- $r_a = \infty$ & $r_b = d$
- $\Delta W = k \frac{q^2}{d}$
- $\Delta W = 8.99 \times 10^9 \times \frac{(1.6 \times 10^{-19})^2}{1.6 \times 10^{-15}} = 2.3 \times 10^{-13}$ J

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

٧٦

Example 06

- Work required to bring two positive charges together.
- What minimum work must be done by an external force to bring a charge $q = 3.00 \mu\text{C}$ from a great distance away (take $r = \infty$) to a point 0.500 m from a charge $Q = 20.0 \mu\text{C}$?



- $\Delta W = -Q(v_b - v_a) = -Q \left(\frac{kq}{r_b} - \frac{kq}{r_a} \right)$
- $r_a = d$ & $r_b = \infty$
- $\Delta W = k \frac{Qq}{d}$
- $\Delta W = 8.99 \times 10^9 \times \frac{3 \times 10^{-6} \times 20 \times 10^{-6}}{0.5} = 1.08 \text{ J}$

34



Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

34

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

Example 07

- Potential above two charges.
- Calculate the electric potential (a) at point A in the figure due to the two charges shown.



- $V_a = VA_2 + VA_1 = k \frac{Q_2}{r_{2A}} + k \frac{Q_1}{r_{1A}}$
- $= 8.99 \times 10^9 \left(\frac{50}{0.3} + \frac{-50}{0.26} \right) = 7.5 \times 10^5 \text{ V}$

34

Dr. Ahmed


Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

34

MCQ

- What is the electric potential at point B?

- 1) $V > 0$
- 2) $V = 0$
- 3) $V < 0$




£1

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

MCQ

- What is the electric potential at point B?

- 1) $V > 0$
- 2) $V = 0$
- 3) $V < 0$




£1

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

MCQ

- Which of these configurations gives $V = 0$ at all points on the

x axis?




£1

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

MCQ

- Which of these configurations gives $V = 0$ at all points on the

x axis?



£1

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II


Potential Due to Any Charge Distribution

- The potential due to an arbitrary charge distribution can be expressed as a sum or integral (if the distribution is continuous):

- $$v = \frac{Q}{4\pi\epsilon_0 r} \quad (v = 0 \text{ @ } r = \infty)$$

- $$v = \sum_{i=1}^n v_i = \frac{1}{4\pi\epsilon_0} \sum_{i=1}^n \frac{q_i}{r_{ia}}$$

- $$v = \frac{1}{4\pi\epsilon_0} \int \frac{dq}{r}$$




ε 0

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

Example 08

- Potential due to a ring of charge.
- A thin circular ring of radius R has a uniformly distributed charge Q . Determine the electric potential at a point P on the axis of the ring a distance x from its center.



ε 1

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II


Potential Due to Any Charge Distribution

- The potential due to an arbitrary charge distribution can be expressed as a sum or integral (if the distribution is continuous):

- $$v = \frac{Q}{4\pi\epsilon_0 r} \quad (v = 0 \text{ @ } r = \infty)$$

- $$v = \sum_{i=1}^n v_i = \frac{1}{4\pi\epsilon_0} \sum_{i=1}^n \frac{q_i}{r_{ia}}$$

- $$v = \frac{1}{4\pi\epsilon_0} \int \frac{dq}{r}$$




- $$v = \frac{1}{4\pi\epsilon_0} \int \frac{dq}{r}$$
- $$= \frac{1}{4\pi\epsilon_0} \int \frac{1}{\sqrt{x^2 + R^2}} \int dq$$
- $$= \frac{1}{4\pi\epsilon_0} \frac{Q}{\sqrt{x^2 + R^2}}$$

ε 0

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

Example 09

- Potential due to a charged disk.
- A thin flat disk, of radius R_0 , has a uniformly distributed charge Q . Determine the potential at a point P on the axis of the disk, a distance x from its center



ε 1

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

$$\bullet A = \pi R_0^2$$

$$\bullet dA = 2\pi R dR$$

$$\bullet \sigma = Q / \pi R_0^2$$

$$\bullet \sigma = dQ / 2\pi R dR$$

$$\bullet \frac{dQ}{Q} = \frac{2\pi R dR}{\pi R_0^2}$$


$$\bullet dQ = \frac{2QR dR}{R_0^2}$$

$$\bullet V = \frac{1}{4\pi\epsilon_0} \int \frac{dQ}{\sqrt{x^2 + R^2}}$$

$$\bullet V = \frac{2Q}{4\pi\epsilon_0 R_0} \int_0^{R_0} \frac{R dR}{\sqrt{x^2 + R^2}}$$

ε 4

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II



$$\bullet V = \frac{Q}{2\pi\epsilon_0 R_0} \int_0^{R_0} \frac{R dR}{\sqrt{x^2 + R^2}}$$


• V

$$= \frac{Q}{2\pi\epsilon_0 R_0} (\sqrt{x^2 + R^2}) \Big|_{R=0}^{R=R_0}$$

$$\bullet V = \frac{Q}{2\pi\epsilon_0 R_0} [\sqrt{x^2 + R_0^2} - x]$$

ε 5

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II



Equipotential Surfaces


• An equipotential is a line or surface over which the potential is constant.

• Electric field lines are perpendicular to equipotential.

• The surface of a conductor is an equipotential

$$\bullet (E_{//} = 0 \rightarrow \partial V / \partial s_{//} = 0)$$


• Equipotential lines (the green dashed lines) between two oppositely charged parallel plates. Note that they are perpendicular to the electric field lines (solid red lines).



ε 6

Dr. Ahmed ElShafee, ACU : Spring 2015, Physics II

• Equipotential lines (green, dashed) are always perpendicular to the electric field lines (solid red) shown here for two equal but oppositely charged particles.



ε 7



Thanks, ...
See you next week (ISA), ...